



## Rehabilitation Protocol: Combined Anterior Cruciate Ligament (ACL) Reconstruction and Meniscal Repair

Name: \_\_\_\_\_

Date: \_\_\_\_\_

Diagnosis: \_\_\_\_\_

Date of Surgery: \_\_\_\_\_

**Phase I (Weeks 0-4)**

- **Weightbearing:** Partial weight bearing (50%) as tolerated with brace locked in full extension
- **Hinged Knee Brace:** Locked in full extension for ambulation and sleeping
- **Range of Motion:** No flexion greater than 90 degrees
- **Therapeutic Exercises**
  - No deep bends for first 4 months
  - Quad/Hamstring sets
  - Heel slides
  - Non-weightbearing stretch of the Gastroc/Soleus
  - Straight-Leg Raise with brace in full extension until quad strength prevents extension lag

**Phase II (Weeks 4-6)**

- **Weightbearing:** As tolerated – discontinue crutch use
- **Hinged Knee Brace:** Discontinue brace use when patient has achieved full extension with no evidence of extension lag
- **Range of Motion:** Maintain full knee extension – work on progressive knee flexion
- **Therapeutic Exercises**
  - No deep bends for first 4 months
  - Closed chain extension exercises
  - Hamstring curls
  - Toe raises
  - Balance exercises
  - Progress to weightbearing stretch of the Gastroc/Soleus
  - Begin use of the stationary bicycle

**Phase III (Weeks 6-16)**

- **Weightbearing:** Full weightbearing
- **Range of Motion:** Full/Painless ROM
- **Therapeutic Exercises**
  - No deep bends for first 4 months
  - Advance closed chain strengthening exercises, proprioception activities
  - Begin use of the Stairmaster/Elliptical
  - **Can Start Straight Ahead Running at 12 Weeks**

**Phase IV (Months 4-6)**

- Gradual return to athletic activity as tolerated
- Maintenance program for strength and endurance

Comments:

Frequency: \_\_\_\_\_ times per week

Duration: \_\_\_\_\_ weeks

Signature: \_\_\_\_\_

Date: \_\_\_\_\_



## Rehabilitation Protocol: Anterior Cruciate Ligament (ACL) Reconstruction with Allograft (Achilles, tibialis anterior, or bone-patellar-bone)

Name: \_\_\_\_\_

Date: \_\_\_\_\_

Diagnosis: \_\_\_\_\_

Date of Surgery: \_\_\_\_\_

**Phase I (Weeks 0-4)**

- **Weightbearing:** As tolerated with crutches (may be modified if concomitant meniscal repair/meniscal transplant or articular cartilage procedure is performed)
- **Hinged Knee Brace:**
  - Locked in full extension for ambulation and sleeping (**Weeks 0-1**)
  - Unlocked for ambulation and removed while sleeping (**Weeks 1-4**)
- **Range of Motion** - AAROM → AROM as tolerated
- **Therapeutic Exercises**
  - Quad/Hamstring sets and heel slides
  - Non-weightbearing stretch of the Gastroc/Soleus
  - Straight-Leg Raise with brace in full extension until quad strength prevents extension lag

**Phase II (Weeks 4-12)**

- **Weightbearing:** As tolerated -- discontinue crutch use
- **Hinged Knee Brace:** Discontinue brace use when patient has achieved full extension with no evidence of extension lag
- **Range of Motion** - Maintain full knee extension - work on progressive knee flexion
- **Therapeutic Exercises**
  - Closed chain extension exercises
  - Hamstring curls
  - Toe raises
  - Balance exercises
  - Progress to weightbearing stretch of the Gastroc/Soleus
  - Begin use of the stationary bicycle

**Phase III (Months 3-10)**

- **Weightbearing:** Full weightbearing
- **Range of Motion** - Full/Painless ROM
- **Therapeutic Exercises**
  - Advance closed chain strengthening exercises, proprioception activities
  - Begin use of the Stairmaster/Elliptical
  - **Can Start Straight Ahead Running at 3 months**

**Phase IV (Months 10-12)**

- Continue with strengthening (quad/hamstring) and flexibility
- Begin cutting exercises and sport-specific drills
- Maintenance program for strength and endurance
- **Return to sports at 10 months**

Comments:

Frequency: \_\_\_\_\_ times per week

Duration: \_\_\_\_\_ weeks

Signature: \_\_\_\_\_

Date: \_\_\_\_\_